

Biological activities and phytochemistry of *Ziziphus mauritiana* from Gwadar (A Review)

SHABANA TAREEN

M.Phil. Scholar, Institute of Biochemistry, University of Balochistan, Quetta, Pakistan

MUHAMMAD ANWAR PANEZAI

Professor, Institute of Biochemistry, University of Balochistan, Quetta, Pakistan

NOOR HASSAN

Professor, Institute of Biochemistry, University of Balochistan, Quetta, Pakistan

JAHANGIR KHAN ACHAKZAI

Department of Natural and Basic Sciences, University of Turbat, Turbat, Pakistan

ABDUL MANAN KAKAR

Institute of Biochemistry, University of Balochistan, Quetta, Pakistan

NAZIMA YOUSAF KHAN

Institute of Biochemistry, University of Balochistan, Quetta, Pakistan

AJAB KHAN TAREEN

Institute of Biochemistry, University of Balochistan, Quetta, Pakistan

ZAHOOR AGHA

M.Phil. Scholar, Institute of Biochemistry University of Balochistan, Quetta, Pakistan

BUSHRA HABIBULLAH

M.Phil. Scholar, Institute of Biochemistry, University of Balochistan, Quetta, Pakistan

NAJMA GUL KHAN

M.Phil. Scholar, Institute of Biochemistry, University of Balochistan, Quetta, Pakistan

Abstract

Plants, being principal source of various medicine performs essential role in the health of the plant and human. *Ziziphus mauritiana* exhibited significant biological activities and have traditional uses for instance Immunostimulant effects, Cardiovascular action, Anti-inflammatory, Antiulcer action, Sweetness repressive, Anti-allergic action, Anti-diarrheal action, hypoglycemic effect, Anti-obese activity, laxative, blood cleansing characteristics, soothing effect, hypnotic, stop vomiting and nausea, in pregnancy for relief of abdominal pain, antidote or a cure towards aconite-poisoning, well Worn vogueish poultices, wounds therapeutic, astrivent, diaphoretic, typhoid fever in children, diarrhea disease, dysentery. The phytochemistry of *Ziziphus mauritiana* revealed the presence of secondary metabolites for instance cardiac glycosides, tannins, polyphenols, alkaloids, pectin, terpenoids, flavonoids, saponin, triterpenoic acids, **phenols**, coumarins, quinones, tarmacs, phytosterols and glycosides. Regarding economic importance the basic raw constituents desired on behalf of pharma harvests, cosmetic industries, flavoring agents plus perfumery products.

Keywords: *Ziziphus mauritiana*, Biological activities, Phytochemistry

INTRODUCTION

In arrange to decide the in vitro flavonoids and complimentary rummaging capacity and resveratrol properties of takes off of *Z. mauritiana*. along with *Z. spina-christi*.

Contrasted through Ascorbic Corrosive. The display thinks about live embraced utilizing in vitro flavonoids strategies such as hydroxyl group, lipid peroxidation, DPPH radical and superoxide radical levelness strategies. Ethanol along with Hexane extricates of *Z. mauritiana* L. as well as *Z. spina-christi* L. takes off were put through the strategies recorded over. The approach about of rutin action uncovered that, Hexane extricate has higher EC50 values than the ethanol extricates of *Z. mauritiana* L. including *Z. spina-christi* L. The elevated free radical rummaging capacity of the plants shown by the lesser EC50 esteem So, the ethanol distangle incorporates a higher flavonoid motion than hexane release. These happen about were contrast with the excellence ascorbic corrosive ordering.

The flavonoids such as cardiac glycosides, saponins, tannins and polyphenols, were illustrious, among others. It is assumed that these vigorous residents acting distinctly or in supportive energy may be reliable for the observed antioxidant deed of these plant types (Abalaka, Mann, & Adeyemo, 2011). *Ziziphus* Process. is the primary critical genres of family Rhamnaceae, this family is largely spread everywhere in biosphere and exploited designed for periods in site pharmaceutical agenda. Greatest theme of this predict was to scouting the biochemical databank, the flavonoids as well as antimicrobial movements of untreated methanolic colliery of *Ziziphus mauritiana* cuff. dissipates out in contradiction of *Enterococcus faecalis* ATCC 29212, *Klebsiella pneumoniae* ATCC 700603, *Streptococcus pneumoniae* ATCC 49619, *Bacillus cereus* ATCC 10876, *Staphylococcus aureus* ATCC 25923, *Escherichia coli* ATCC 10536, *Proteus mirabilis* ATCC 14153 *Pseudomonas aeruginosa* ATCC 27853, also a pathogenic creature; *Candida albicans*.

Originates about phytochemical screening have seem like, the plant excerpt topographies a distinguishable volume of biotransformations components; flavonoids, tannins, phenols, coumarins, cardiac glycosides, quinones, tarmacs, saponins, phytosterols as well as terpenoids. originates about of flavonoids endeavor of leaves mine have looked particularly, IC50 (semi utmost discretion flock) repute of leaves colliery was 0.875 mg/mL, suggested particularly, rescue of plant encompasses a prominent quantity of acceptable radical's accomplishment measure. Extricate of plant has too performed acute level of antibacterial measure in contradiction of *B. cereus* ATCC 10876 (area of hindrance expanse across: 12 mm, MIC: 625 µg/mL, MBC: 1250 µg/mL), *S. aureus* ATCC 25923 (region based on hindrance extent: 12 mm, MIC; 625 µg/mL, MBC: 1250 µg/mL), *S. pneumoniae* ATCC 49619 (sector about curb range: 10 mm, MIC; 156.25 µg/mL, MBC: 312.5 µg/mL) plus antimycotic execution at the side of *Candida albicans* (part of curb extensiveness: 10 mm, MIC: 125 µg/mL). Hence, major methanol citation of *ziziphus Mauritiana* cuff. proceeds off may remain exploited designed for the medicinal of changed humanoid infections. A spacious quantity of floras is consumed via fables treaties voguish India used for handling of lesions, injuries and burns.

Precisely, the disintegrates of *Ziziphus mauritiana* Lam. (Family: Rhamnaceae) stayed skilled on injuries used for recovering. The datum of the exhibition inquiry was to evaluation in vivo wound improving tolerability of systematized topical narrative of ethanoic extract achievement 5% w/w and liquefied extract control 5% w/w and 0.2% w/w of nitrofurazone supervision on drawing out wound suggestion Wistar light sensitive somebody rats. impression which approved by action, in rapports of wound tightening competency, wound terminating, condense in surface zone of injuries,

tissue recapture at the injury section in healed Wistar whitish sensitive one rats give the impression that improvement of epithelial tissue growths angiogenesis, augmentation of sturdy connective tissue allocated to management by *Ziziphus mauritiana*.

Commanding venomous superiority contemplates exposed the non-venomous atmosphere of *Z. mauritiana*. Ethanolic excerpt accomplishment (5% w/w) of *Ziziphus mauritiana* presented 99% wound concreteness on day 16th. These initiates about were furthermore analogous to individuals of an ordinary serene, nitrofurazone. Fascinating definition through ethanolic extract 5% w/w innovative wound insistence plus condensed the injury finishing period, therefore rise in pliable superiority also wound density seems the injury sewing latent of *Ziziphus mauritiana*. In this procedure, the demonstration considers provisions the logical foundation aimed at the conventional usage of plant *Ziziphus mauritiana* privileged the administration of injuries. Although a *Ziziphus mauritiana* has been observed on behalf of its anticancer, antioxidant, antidiabetic as well as nutraceutical actions (Koley et al. 2011), some species, like *Z. mauritiana* occur almost in every region. It is a pattern of a tremendously scarcity-resistant species in addition is a main constituent of the expected flora of the India and Pakistan deserts. In *Ziziphus* diversities have an extensive ritual of collection and farming in both China plus India, which stemmed in enhanced known as well as more extensively researched variabilities than those in additional areas (Cherry, 1985).

Plants are natural pool of antimicrobial representatives of therapeutic values. These representatives are assumed to be virtually free from lateral properties usually related with artificial antimicrobials. So several of the existing day medications are identified to have been secluded from expected causes plus their segregations were centered on the material about the expenditures of the mediators in folklore medication.

PHYTOCHEMISTRY

Numerous complexes such as lipids, triterpenoic acids, alkaloids, pectin A, and glycosides are existing in *Z. mauritiana*. Compound Pectin A have 2, 3, 6-tri-o-acetyl D lactose components, pectin partakes Several therapeutic possessions for example lowering plasma cholesterol binding bile acid, plus anti diarrhoeal (Dahiru et al., 2006). saponins secluded against seeds of *Z. mauritiana* comprise of jujubosides A1, B1, A, B, C as well as acetyljujuboside B. Protojujubosides A, B as well as B1 plus ziziphin exist correspondingly secluded commencing dried out leaves. Jujubosides pledge to partake sedative plus anxiolytic haemolytic, as well as sweetness constraining possessions. Jujuboside A, is recognized as a uncompetitive inhibitor regarding a calmodulin, whichever is a calcium-obligatory protein, ubiquitous which can fix plus standardize an assembly based on various protein purposes, in that way heartrending various diverse cellular determinations for example, muscle contraction, short-term inflammation, metabolism as well as long-term memory, plus immune response (Yoshikawa K et al, 1991; Matsuda H et al, 1999; Kurihara Y et al, 1988; Sharma SC, et al, 1982; Oda K, Matsuda H et al, 2000; Shou C, Feng Z et al, 2002; Zhou Y, Li Y, Wang Z et al, 1994) *Ziziphus mauritiana* has triterpenoic acids for example colubrinic acid, 3-O-cis-p-coumaroylalphitolic acid, 3-O-cis-p-oumaroylmaslinic acid, betulonic acid, oleanonic acid, 3-O-trans-p-coumaroylmaslinic acid, oleanolic acid, 3-O-trans-p-

coumaroylaliphitic acid, aliphitic acerbic, zizyberenic acid plus betulinic acid. oleanolic acid in addition to Betulinic acid possess validated discerning cytotoxicity in contrast to a numeral about exact cancer groups, moreover betulinic acerbic. or acid possess anti-inflammatory action in addition to antibacterial assets. (Lee S, Min B et al, 2003) (Kundu AD et al, 1989).

PLANT PROFILE

leaves of ziziphus mauritiana are alternate, complete, by means of three distended veins of rosenthal, then 2 - 7 cm means 0.79 - 2.8 in long; numerous species remain ephemeral, some more illustrious flowers of ziziphus Mauritian are small, unobtrusive yellow-green. fruit of this plant is comestible drupe, red, or black, yellow-brown, globose or rhombus, 1 to 5 cm (0.39 - 2.0 in) elongated, frequently extremely sweetened as well as syrupy, momentous about a period in stability plus flavor. altitude of Ziziphus mauritiana tree varies concerning the place. It is 6-9 m high in many countries whereas in additional countries it is 9-12 m high. The trees of ziziphus mauritiana are additional branched than others. Their fruit size is also diverse concerning the agriculture plus environmental circumstances (Ji X, Peng Q, Yuan Y, et al, 2017).



Figure Error! No text of specified style in document..1 whole plant of ziziphus mauritianass

SPREADING PLUS EXISTENCE

Ziziphus Refine is the most significant genera of the Rhamnaceae family, which is broadly scattered in the entire biosphere and used in medication systems for centuries. *Z. mauritiana* is one of the common specie in 40 species of ziziphus which are recorded till present day.

The best recognized species of this plant is *Z. zizyphus* (Jujube). Further species comprise *Z. lotus* from the Mediterranean, *Z. spina-christi* in southwestern Asia, as well as Ber (*Z. mauritiana*) is found in western Africa in the direction of India and in the Caatinga region of Brazil *Z. joazeiro* cultivates. Many species harsh occur on every mainland for example *Z. mauritiana* Lam. plus *Z. spinachristi* (L.). the fruits of *Z. mauritiana* plus *Z. spina-christi* are very nutritious.

The focal locality of ziziphus mauritiana includes Meditarranean, America in addition to Asia areas (tropical and subtropical). As the plant of ziziphus mauritiana is instinctive to china so china is accountable for 90% manufacture of this plant in the

world. *Ziziphus mauritiana* variations includes: Illaichi, Gola, Umran, Banarasi Karaka, Seb, Bawal-Sel-1, Thar Sevika, Sanaur-2 Ponda Tikadi, Katha, Thar Bhuhhraj, etc.

TAXONOMY AND NOMENCLATURE:

Taxonomical classification

Kingdom! Plantae
Division! Magnoliophyta
Class! Magnoliopsida
Order! Rosales
Family! Rhamnaceae
Genus! *Ziziphus*
Species! *Mauritiana*

Taxonomic Description

Ziziphus mauritiana belongs to family Rhamnaceae and recognized as jujube tree (Morton, 1987; Michel, 2002). *Mauritiana* is the specie of genus *Ziziphus*. *Ziziphus* is a member of Rhamnaceae family which named afterwards the genus *Ramus*. Fruits of Rhamnaceae family are dry or remain drupes plus narrowly associated to surplus family recognized as vitaceae, the species of this family embraces major economic importance having are barriers as a fruit. *Ziziphus* name is associated headed for a Greeks word plus Arabic ziziphon used for the jujube or *mauritiana*. Around two main tamed jujubes, *Ziziphus mauritiana*. ber plus *Ziziphus jujube*. cultivation of these dual species are spread concluded a huge regions of the world. These species have an extensive variety of morphological difference commencing shrubs to a medium or small sized trees in which some are erect, spreading or else semi-erect. The Height of *ziziphus mauritiana* differ from 3-4m to 10-16m some rare trees have 20m long.

Ziziphus Mauritiana plant is in various instances predictable as whuya in Kilba (Nigeria) and magarya in Hause.

There are nearly 86 species of genus *ziziphus* (Azam-Ali et al., 2006) of small trees or prickly shrubs that are dispersed in subtropical areas to hot-temperate sections of the world. In india all over the place 20 species regarding *ziziphus* are existing (Awasthi & More, 2009) plus *ziziphus* 14 species existing in China (Zhao & Liu, 2009). Amid them, the two species of *ziziphus* family are *Ziziphus mauritiana* that is also identified by means of Indian jujube or else ber is an extra tropical as well as immortal species in subUnit *Perdurans* plus *Ziziphus jujuba* which is known by means of Chinese jujube then Chinese date is an additional cold-tough as well as ephemeral species trendy Sector *Ziziphus*, be situated versatile trees cultured broadly a small cultural yields on behalf of fruits (Azam-Ali et al., 2006; Pareek & Yahia, 2013).

Ziziphus mauritiana has around other relatively to 170 cultivars in region of India country. (Azam-Ali et al., 2006) as well as china has above 750 *Ziziphus. jujuba* cultivars. (Zhao & Liu, 2009), that is differ broadly in tree habit, shape regarding leaf, form of fruit, color, flavour, size, possession eminence plus fruiting period. *Z. mauritiana* is different from *z. jujuba* be ause of the thick silky underneath of its leaves (Morton, 1987). If the commercial as well as wild species are hybridized then in resulting an intermediate form are formed (Hocking, 1993). In the country of Australia

dual natural *Ziziphus* species remain existing *Z. quadrilocularis* F. Muell. plus *Z. oenoplia* (L.) Mill. (Grice, 2002), mutually in Division Perdurans. *Z. oenoplia* is as well inherent towards Sri Lanka as well as tropical Asia in addition to India (Grice, 2002).

Local Name

Z. mauritiana a member of Rhamnaceae family The Plant of *ziziphus mauritiana* is known by several names. The local name of *Ziziphus mauritiana* is Qunar. Its common name Ber. In Sans it is known as Ajapriya, badara, madhuraphala karkandhu, kuvala, in Hindi it is called Baer, ber in Bengali it is Kool, ber, boroi in Marathi language it is known by Bar, bera in Gujarati it is known by Bor, bordi in Telgu *z. mauritiana* known by Reegu, karakandharu gangareegu in Tamil. *z. mauritiana* called by Elandai, yellande elladu in Kanda language this plant known by Yalachi, elanji, in Malalum. known by Eentha in Oriya *z. mauritiana* is recognized by Barkoli, bodori, bodokoi, in Rajasthan this plant is familiar by Bor, bordi. In Sanskrit it is called by Rajabadari name in Punjabi language *z. mauritiana* is familiar as Beri. In Assamese it is recognized by Bogori. In Malayam it termed as Badaram. In Kannada language this plant called by Bogari. In Tamil *z. mauritiana* known by Vadari name. In Telugu it is familiar as Renu. In Urdu it is recognized by Ber. While in sindi *z. mauritiana* is known Jangri.



Figure Error! No text of specified style in document..2 leaves of *ziziphus mauritiana*

TRADITIONAL USE

Round about 60 percent population of the world use Traditional medicines. Not only a ruler areas used *z. mauritiana* for primary fitness repair but developing countries also use it. Modern medicines are mostly used in developed Countries although traditional drugs are consequent from minerals, organic matter, and medicinal plants that are why herbal treatments remain only organized commencing therapeutic florae. Usage about florae by means of an inventor of medication has been inherited from years and it is a significant constituent of healthiness treatment arrangement in Pakistan plus india. The arrangements of medication in Pakistan, greatest general practitioner formulate plus dispense their peculiar methods; hence this necessitates a proper credential as well as research. Government, academic and public interest is rising exponentially in traditional drugs by reason of improved prevalence of adversative medication responses as well as commercial burden as regards the present scheme about medication.

In Khandesh area seven contributors about the family as regards Rhamnaceae happened specifically North Maharashtra Section. Rhamnaceae family has specie Jujuba which is being inherent to Chinese medication since 4000 years. Furthermore, it is a portion of Iranian traditional medication as it owns laxative plus blood cleansing characteristics (Najafi, S., Sanadgol, et al, 2010). The fruits are testified to have a soothing effect as well as are suggested as a hypnotic. *Z. mauritiana* is used to stop vomiting and nausea. It also prescribed in pregnancy for relief of abdominal pain. *Z. mauritiana* is furthermore Specified by way of antidote or a cure towards aconite-poisoning, plus well Worn vogueish poultices as well as in other preparations of wounds therapeutic. As an astrinve with catechu the leaves of *Z. Mauritiana* are eaten. They are considered as diaphoretic plus approved for typhoid fever in children. Broth as regards the bark of *ziziphus mauritiana* remains prescribed on behalf of management as concerns diarrhea disease plus dysentery. (Peto Azam-Ali Set al, 2006) (Anonymous. 1989) (Awasthi OP, 2009) (Pareek OP 2001) (Bhandari MM 1978) (Bhattacharjee SK 2004).

ECONOMICAL USE

Aromatic plus medicinal florae are Considered as economically significant plants for the reason that these stays the main cause as regards basic raw constituents desired on behalf of pharma harvests, cosmetic industries, flavoring agents plus perfumery products, therefore they contribute obsessed by GDP of the country through earning of overseas currency (Aiyelaagbe,2001 and Augusti, K.T, 1996). There are several variations are reported various scientists have reported variation among ber cultivars. in Pakistan the Botanical description of local gene group has not taken seriously up as well as records of these local cultivars are commonly missing, whereas, its better product varieties are identified by its care-free creation demands for low wages masses. Thus the manipulation of such assortments may enhanced the commercial yielding of ber (*Z. mauritiana*) in Pakistan with lesser inputs, better produce in addition to improved nutrition

REVIEW LITERATURE

Ziziphus mauritiana as well as *Z. jujuba* remain species as regards genus *Ziziphus* Tourn. Which goes towards Rhamnaceae Family. *Ziziphus* is that name which relates to a word as regards Arabic language cast-off alongside North African coastline, the word *zizoufo* castoff on behalf of *Z. lotus* (L.) Desf., then also connected through words as regards earliest Persian language known as *zizum* likewise *zizafun*; whereas in prehistoric Greeks language the term as regards *ziziphon* used on behalf of *Z. mauritiana* (Peto Azam-Ali et al 2006). Prehistoric of chalcolithic age eat the Jujubes and *Z. mauritiana* (1500-1000 B.C.), as well as in both countries India plus China the fruits of *Z. mauritiana* have been farming since 400 years. These fruits of *ziziphus* have been relatively often mentioned in epics, medical texts, Yajurveda, Sutras, along with this the additional literature determined via researchers for instance Panini, Patanjali plus Kautilya, 3 categories of plant jujubes remain cited in Brahmanas plus Yajurveda: Koal either Ktmila possesses usual sized fruit, the fruit of Badara possess bIg Size, as well as Karkandhu possess orange-reddish colored either brown fruit; while *Ziziphus*

mauritiana appeared as the first two types, the third kind, possessing red colored fruit, seem like elect *Z. nummularia* (Anonymous 1989).

The plant of *z. mauritiana* possess fruit of extremely dearth hardy plus native in India. In india specially in thar desert It is considered as a dominant constituent of the natural flora. It is successfully cultured unexpectedly in greatest bordering environment about the regions as regards torrid plus subtorrid .in india the improved ber (*z. mauritiana*) have 90,000 plantations, that possess average as regards 8.34 tons' productivity for every hectare (Awasthi OP 2009). Though, entirely jujubes persist moderately trivial harvests even though claim on behalf of manufacture persist fixed in numerous Zones as concern. India wherever they remained formerly acclimatized. The plant of *Ziziphus mauritiana* be there measured by means of an under-utilized yield, in addition to this it is involved in nationwide programme on under-utilised yields (Pareek OP 2001). Regardless about the truth a particular *Z. mauritiana* owing fabulous pharmaceutical possessions, accredited via a various cluster of subsidiary metabolites on behalf of instance alkaloids, pectin, terpenoids, flavonoids as well as saponin the plant of *Z. mauritiana* is not either measured by means of neither vital therapeutic floras nor adventure on behalf of therapeutic expenditure in central succession therapeutics. Life and ailments go together the apartment there is life, diseases are assured to exist. Addiction and sustainability of gent and animal routines has been rotating around plant lifetime through their usages as nutrition, fibers and accommodation, however also vegetation have been cast-off to handle and affluence diseases, consequently, the usage of plant lifetime a s drug handling is an earliest and trustworthy implementation (Arshad and Rao, 2001). Therapeutic plant life shows a significant part in human fitness maintenance. Around 80% of world inhabitants be contingent on consumption of the typical medication, which is commonly constructed on herbal considerable (WHO, 1993).

Methodical investigation easily reached on a brilliant range of therapeutic flowers point out that propitious phytochemicals be able to established for several fitness matters (Gupta, 1994). Leaves of *Ziziphus Mauritiana* are used in medication of pustules, woundsand, inflammation, diarrhea, gonorrhoea (Michel, 2002). Their leaves are furthermore expenditure in the medication of liver infections, fever in addition to bronchial asthma (Morton, 1987). Compounds of Antioxidants are excellent complexes that can stay, intersect or prevent the association of oxidative responses through counterbalancing open radicals by means of endowment of hydrogen electron or atom, slaking singlet plus triplet oxygen then chelating metals also for that purpose play a positive role on the way to cultivating shelf-life of nutrition harvests as pleasantly as dropping the presence of various sicknesses for instance growing older, inflammation and cancer (Lobo et al., 2010; Dalleau et al., 2013; Khadri et al., 2010).

At this time, due to the professed poisonous consequences of manmade antioxidants and further than single microbial medicine opposition (MDR), the usage of floras consequent customary antioxidants and bioactive, is consequently center of consideration of existing scientific investigation (Alnajjar et al., 2012).Vegetation mostly constructed phenolic antioxidant compounds, owing in the direction of its exclusive organizational structures in addition a pair of biological movements, perform an auspicious role in the cure of positive transmissible diseases and cancer (Miyasaki et al., 2013; Prasad et al., 2012; Bhalodia as well as Shukla, 2011). Allocated on the way to the revitalization about upheaval vogueish the payments as regards herbal lifetime by

means of a base as concerns drug plus meals, presently scientists are commonly affianced in discovering plus estimating bioactive conformation plus organic concepts of supplementary also better vegetation (Elless et al., 2000; Shahidi, 2009; Lucock, Liu et al., 2008 ,2004;). (*Z. mauritiana*) an associate since the local Rhamnaceae family by way of neighborhood recognize Ber, a fruitlet tree cultivating in a steamy and sub-steamy constituencies of the biosphere. Various portions of *Ziziphus mauritiana* tree have stood castoff in the mutual medicinal remedy for the administration of any of a kind disorder such as depression, asthma, ulcers and allergies, (Marwat et al., 2009). Conversely, approximately lessons so enthralled take moreover examined the phenolic arrangement of fruitlet of the *Z. mauritiana* plus clarified the methodical source designed for the specific portions of *Z. mauritiana* herb for the administration of a diabetic disease, infection also cancer disease (ulcer) (Bhatia as well as Mishra, 2010; Memon et al. 2000; Gupta et al., 2012; Cisse et al., 2012; Siddharth et al., 2010).

The Leaves about plant as regards *Z. mauritiana* applied vogueish the vicinity to sores, in addition the roots remain used to management and end skin disorders (Adzu et al., 2001). Natural products containing plants as well as vegetables are used as primary cause for the improvement of drug to treat various diseases ever since prehistoric time. The plant of *Z. Mauritiana* is traditionally manipulate for medicinal dedications since centuries. *Z. Mauritiana* holds much rank worldwide for the reason of its medicinal profits including pharmacological features and because of this the farming of many species of herb takes place in various countries of the world. Modern studies related to the plant of *ziziphus mauritiana* illustrate that the starting point of this herb is Africa as well as Australia.

The nutritive composition of *z. Mauritiana* plant comprises of amino acids, carbohydrates, oil, proteins, vitamins in addition to phenolic compounds (Elaloui M, Laamouri A et al,2015). The other elements calcium, terpinoids, iron, flavonoids, glycosides and phosphireare also present in *z. Mauritiana* specie of plant. Along secondary metabolites the Bioactive compounds are also present for instance oleic acid in addition to palmitic acid (Hernandez F, Noguera-Artiaga L, et al, 2016). This plant has the capability to control the aging progress in women demonstrated by new studies. *Z. Mairitiana* also used for treatment of numerous forms of ulcer likewise intestinal as well as rectal cancer. vitamin C, terpenoids, phenolic compounds in addition to polysaccharides are considered as a healthy constituents of fruit of *z. Mauritiana* (Gao QH, Wu CS, et al, 2013) every part of *z. Mauritiana* have different talents for instance leaves possess therapeutic properties, fruit possess anti-bacterial as well as antifungal capabilities and seeds with oil possess antimicrobial respectively.

Approximately pharmaceutical florae Possess antibacterial action, antioxidant action then antifungal actions consequently they might remain cast-off for handling about numerous humanoid disorders as therapeutic agents for example to treat the acne, curing as concerns injuries, boil plus numerous skin(membrane) infections, subsequently plants possess excessive prominence vogueish area the medication organization (Arora, D et al, 1999). Pharmaceutical representatives on behalf of treatment about numerous contagious syndromes in people discovered from the time when long previously the attentiveness as regards people over the existence as concerns microorganisms (Sofowora, A., 1982).

The arrival about antibiotic opposition now pathogenic microorganisms be there the largest dare on behalf of industry of healthcare in mutually established as

well as emerging countries. advent regarding MDR (multidrug resistant) microorganism bacteria be present the major dare along with encumbrance in hospital plus community sceneries (Cosgrove, S.E et al, 2003). Currently, the organization of healthcare remains further progressive vogueish antibiotic cure, despite the fact difficulties about transmittable ailments remain the greatest imperative origins of impermanence vogueish patients which are hospitalized. Artificial antibiotics possess several adjacent possessions that might persist permanent now cured victims. For the removing of this difficulty, those movements necessarily be occupied on behalf of monitoring as well as decreasing as regards such gears then it may be completed via monitoring usage regarding antibiotics, via sympathetic the heredity character about antibiotic confrontation, emerging of fresh antibiotics plus altering the pharmaceutical policies (Nascimento, G.G.F. et al, 2000).

A diversity of bioactive ingredients of known healing values are produced by a lot of herbal plants. these medicinal plants possessed an Antimicrobial activity which has one of the greatest important therapeutic values (. Bolla, J.M., Alibert-Franco, et al, 211). The composites, which can either, stop or destroy the pathogens minimum or no poisonousness towards mass cells remains observed by way of greatest applicants on behalf of emerging modern antimicrobial supervisors (Qureshi, K.A., et al, 2016; Alanazi, A.S., et al, 2016; Lee, C.K., Kim, et al, 1998). About antimicrobial activities Many information are published all over the world in the previous few years which recommended that this activity of therapeutic flora have been amplified (Grosvenor, P.W., Supriono, et al, 1995; Ahmad, I., Beg, et al, 2001; Abu-Shanab, B.A., Adwan, et al, 2008).

Ziziphus which remains the supreme significant genera regarding family Rhamnaceae, it exists broadly scattered all over the world plus it remains cast-off as locale medication systems from centuries. Up till then, nearby 40 species of genera of Rosacea family (Ziziphus) are recorded and in this record Ziziphus. mauritiana is very mutual, specifically in dry areas.the species of Ziziphus are very good source of vitamins, fibers, carbohydrates and proteins (Parmar, P., Bhatt, et al,2012). the leaves of Ziziphus mauritiana remain extremely generous in ore, therefore approximately faunae feed on leaves of z. mauritiana to get profit for their healthiness for example cattle, goats, camels etc.

(Morton, J.et al, 1987). The fruits of Z. mauritiana exist extremely generous in filaments as well as own purgative action along with thus, much advantageous on behalf of fitness about human beings (Parmar, P., Bhatt, et al,2012). Leaves of Z. mauritiana Lam. Used to treat many diseases like asthma, fever and liver problem, while the precipitous of extract of z. mauritiana root is the treatment for therapeutic of wounds (Dahiru, D., Sini, et al, 2006).

BIOLOGICAL ACTIONS OF LEAVES REGARDING ZIZIPHUS MAURITIANA

Immunostimulant effects

The extraction of leaves concerning Z. Mauritiana remained establish in the direction of excite the chemo tactic, phagocytic as well as intracellular slaughter potency of neutrophils regarding human beings “infection struggling by means of WBCs” at 5 -50 micro g/ml.

Cardiovascular action

Neo-lignan which is secluded commencing leaves regarding mauritiana has ability to expand or rise the discharge concerning endogenous hormone prostaglandin I₂ “best persuasive usual interdictor as regards platelet accumulation however discovered as well as an authoritative vasodilator” commencing rat aorta via equal near 25.3 % next to 3 micro g/ml.

Anti-inflammatory

Excerpt regarding Ziziphus mauritiana possess important anti-inflammatory action in contrast to carrageenan-persuaded concerning rat appendage edema. Percentage of curbing about appendage edema proceeding three hours subsequently carrageenan organization formed via the excerpt of leaves concerning Z. mauritiana on the dosage as regards 200, 600 plus 400 mg/kg persisted 44.5%, 81.8% plus 62.2% respectively once compared to regulate.

Antiulcer action

Z. Mauritiana leaves extract owns momentous as well as dosage contingent antiulcer action via pylorus ligation representation rats. antiulcer action of z. mauritiana extract can be accredited to the aforementioned cytoprotective as well as Anti-secretory deed.

Sweetness repressive

Out of possession of the leaves excerpt regarding z. Mauritiana inhibitors of Triterpenoid sweetness were sequestered. Leaves extract concerning Z. mauritiana remain set up in the direction of overpower sugary sensitivity sense vogueish fly “Pharmaceutics”, hamster plus in rat. Anti-sugary ingredients which are separated from Z. mauritiana comprised jujubosaponins II, III, IV, V as well as VI in addition as of the leaves, ziziphus saponins I -III after dehydrated fruit as well as jujuboside B commencing the seeds as well as leaves. Ziziphin plus jujubosaponins II and III, just three antisweet saponins from z. mauritiana herb by means of acyl groups, remained able to four periods extra energetic now conquering the sugary flavor regarding sucrose compound as compared towards other anti -sugary ingredients besides in that way decreasing obesity in overweight otherwise diabetic human beings.

Anti-allergic action

In aqueous extract concerning z. mauritiana leaves has the anti-allergic action this remained deliberate via determining its repressing consequence proceeding hyaluronidase “depress testes” stimulation vogueish vitro. Ziziphus mauritiana be there exposed to possess threatening anti-allergic action.

Anti-diarrheal action

The extract of z. Mauritiana leaves in the experiment of castor oil-induced diarrhea produced an important inhibitory action in contrast to castor lubricant plus Mgso₄ persuaded diarrhea vogueish rats.

Hypoglycemic effect

examination variance results designated momentous decrease (P = 0.001) regarding glucose–triglyceride–cholesterol as well as very low density lipoprotein levels vogueish

cluster 3 vogueish contrast per bunch 2. *Ziziphus mauritiana* correspondingly improved High density lipoprotein stages ominously ($P=0.001$) vogueish contrast by set 2. As well, the excerpt Decrease low density lipoprotein level of diabetic rats, however this one is not momentous ($P=0.12$) in appraisal with set 2.

Anti-obese activity

Alcoholic extract regarding leaves of *Z. Mauritiana* indicated anti-obese possessions by reducing the weight of body, food consumption, serum glucose as well as lipid levels plus internal organs in addition to fat pack loads in nutritional obese or fat rats. Conclusion which formed by it remained equivalent thru such formed via Sibutramine which is a standard anti-obese medicine.

Leaves used as medicine

Plenteously of therapeutic plants are existed in whole world. *Z. mauritiana* is Individual of them. It is the genus of Rhamnaceae family. 40 species of *Z. Mauritiana* plant has been discovered with valuable uses scattered worldwide but in these species only 4 species are designed for medicinal determinations. This plant of *Z. Mauritiana* is intended for improvement of medicines in various countries including India, China, in addition to several Asian countries (Stadlmayr B, et al, 2013) color regarding leaves of *Z. mauritiana* plant is green and having yellowish flowers. Therefore, the basic methanolic excerpt of *Z. mauritiana* Lam. incomes off may be applied intended for the medicinal of various human contagions. A spacious number of plants remain consumed by fables conventions in India country for handling of cuts, injuries also burns.

Specially, dissipates out of the *Ziziphus mauritiana* Lam. This Family (Rhamnaceae) trained on lesions designed for recovering. Leaves of this plant stay used in the medication of abscesses, woundsand, swelling, diarrhoea, gonorrhoea (Michel, 2002). Family (Rhamnaceae) are furthermore wield in medication of fever, bronchial asthma plus liver sicknesses (Morton, 1987). Even though in Asian countries the Specie of *Ziziphus* of Rhamanceae family has been extensively and mostly used, mostly in Taiwan as well as in China for handling of allergies, chronic bronchitis, insomnia, depression, constipation, urinary problems plus liver diseases (Li et al. 2005).

Many diseases for example diabetes, cancer, skin diseases and liver infections are treated through this plant. *Z. Mauritiana* is also intended for cleansing of blood. Various forms of fruits plus vegetables have chemoprotective properties towards the destruction triggered by anticancer medicines to organs as well as tissues. Numerous plants contain natural plus bioactive complexes which are used in the improvement of new medicines for instance anticancer medicines (Lozoya, M.,et al, 1989) The nutritional conformation of it's also contains phenolic composites which makes it able to treat various types of diseases (Aiyelaagbe, O., et al, 2001).

Various parts of *Z. mauritiana* plant have diverse abilities for example leaves having healing possessions, fruit having anti-bacterial plus antifungal capabilities as well as seeds with oil having antimicrobial respectively. the stem of *Z. Mauritiana* used in the form of powder in Some countries, while powder of bark as well as leaves of *Z. Mauritiana* use for the management of lesions. As this herb have presented to be the best basis of polyphenolic complexes, it is necessary for patients of diabetes in order to regulate the diabetes.

NUTRITIONAL VALUE

The leaves plus fruits of ber plant (*Z. mauritiana*) from dietary estimation are very necessary for being flusher in inanimate ingredients vitamin C along with other crucial nutritional constituents. The species of *Ziziphus* are very amusing source of carbohydrates, vitamins, fibers and proteins. As leaves of ber plant are opulent in natural resources, that's why many faunae like cattle, goats, camels nurture on these shrubberies to attain improvement for their power fruitlets of ber plant are very gorgeous in filaments plus keep purgative action and thus, much advantageous for human healthiness. Many main minerals for instance potassium, calcium in addition to phosphorus, other minerals for example iron, copper plus zinc are also present in fruits of *Z. Mauritiana*. (Karthikeyan, A., Shanthi, et al, 2009).

Different vitamins comprising vitamin B plus C are also existing in gre. comat volumes. The *Z. mauritiana* leaves extract contains of amino acids, proteins in addition of various fatty acids comprehensive of oleic, palmitic, palmitoleic and linoleic acids. dietary fibers and some fructose insides also present in This plant which control sugar levels of blood. Various types of sugars such as rhamnose, sorbitol, sucrose and glucose are also the fragment of its nutritive configuration. Both leaves as well as fruits of *Z. mauritiana* are very vital from nutritional viewpoint for being wealthier in mineral innards (Ivanova, 1982) vitamin C (Wang et al., 1992) plus other significant nutritional contents.

REFERENCES

1. Abalaka, M., Mann, A., & Adeyemo, S. (2011). Studies on in-vitro antioxidant and free radical scavenging potential and phytochemical screening of leaves of *Ziziphus mauritiana* L. and *Ziziphus spinachristi* L. compared with Ascorbic acid. *Journal of Medical Genetics and Genomics*, Vol. 3(2), pp. 28 - 34.
2. Abu-Shanab, B.A., Adwan, G., Adwan, K.M., Abu-Shanab, F., Efficacy of aqueous and ethanol extracts of some Palestinian medicinal plants for potential antibacterial activity, *Islam. Univ. J. Gaza*, 2008; 16(2): 7786.
3. Adebooye OC, Opabode JT (2004). Status of conservation of the indigenous leaf vegetables and fruits of Africa. *Afr. J. Biotechnol.* 3: 700-705.
4. Adzu B, Amos S, Wambebe C, Gamaniel K (2001). Antinociceptive activity of *Ziziphus spinachristi* root bark extract. *Fitoterapia*, 72: 334350
5. Ahmad, I., Beg, A.Z., Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens, *J. Ethnopharmacol*, 2001; 74(2): 113-23.
6. Aiyelaagbe, O., Antibacterial activity of *Jatropha multifida* roots, *J. Fit.*, 2001; 72(5): 544-546.
7. Alanazi, A.S., Qureshi, K.A., Elhassan, G.O., Elsayed El-Agamy E.I., Isolation, purification and characterization of antimicrobial agent antagonistic to *Escherichia coli* ATCC 10536 produced by *Bacillus pumilus* SAFR-032 isolated from the soil of Unaizah, Al Qassim province of Saudi Arabia, *Pak. J. Bio. Sci.* 2016; 19(5): 191-201.
8. Ali Al Ghasham, M. A.-A. (2017). Phytochemical Screening, Antioxidant and Antimicrobial Activities of Methanolic Extract of *Ziziphus mauritiana* Lam. Leaves Collected from Unaizah, Saudi Arabia . *International Journal of Pharmaceutical Research & Allied Sciences*, 6(3):33-46 .
9. Alnajar, Z.A.A., M. Abdulla, M.H. Ali, A.M. Alshawsh and Hadi. 2012. Acute toxicity evaluation, antibacterial, antioxidant and immunomodulatory effects of *Melastomamalabathricum*. *Molecules*, 17(3):3547-3559.
10. Anonymous. The Wealth of India Raw material Vol XI X-Z, Council of Industrial and Scientific Research, New Delhi, 1989. p.111-24
11. Anonymous. The Wealth of India Raw material Vol XI X-Z, Council of Industrial and Scientific Research, New Delhi, 1989. p.111-24
12. Arora, D., Kaur, J., Antimicrobial activity of spices, *Int. J. Antimicrob. Agents*, 1999; 12 (3): 257-262.
13. Arshad, M. and A. Rao, 2001. Medicinal Plants of Cholistan Desert. In: R. Anwar, Haq, N., Masood, S., Medicinal Plants of Pakistan, P:1.
14. Augusti, K.T., Therapeutic values of onion and garlic, *Indian J. Exp. Biol.*, 1996; 34(3): 634-640.

15. Awasthi O.P., More T.A. (2009) Genetic diversity and status of *Ziziphus* in India. *ISHS Acta Horticulturae*, 840, 33–40.
16. Awasthi OP, More TA. Genetic diversity and status of *Ziziphus* in India. *Acta Hort.* 2009; 840:33-40.
17. Ayodele AE (2005). The medicinally important leafy vegetables of south Western Nigeria. *Ethnobotanical leaflets*. Web journal <http://www.siu.edu/~eb/leaflets/ayodele.htm>.
18. Azam-Ali S., Bonkougou E., Bowe C., DeKock C., Godara A., Williams J.T. (2006) *Fruits for the Future 2*. Revised edn. Ber and other jujubes. Southampton, UK: University of Southampton, International Centre for Underutilized Crops. 289 pp.
19. Bhalodia, N.R. and V.J. Shukla. 2011 Antibacterial and antifungal activities from leaf extracts of *Cassia fistula* L. An ethnomedicinal plant. *Journal of Advanced Pharmaceutical Technology and Research*, 2: 104-109
20. Bhandari MM. *Flora of the Indian Desert*, Scientific Publishers, Jodhpur, India, 1978. p.114-5.
21. Bhatia, A. and T. Mishra. 2010. Hypoglycemic activity of *Ziziphus mauritiana* aqueous ethanol seed extract in alloxan induced diabetic mice. *Pharmaceutical Biology*, 48: 604610.
22. Bhatia, A. and T. Mishra. 2010. Hypoglycemic activity of *Ziziphus mauritiana* aqueous ethanol seed extract in alloxan induced diabetic mice. *Pharmaceutical Biology*, 48: 604610
23. Bhattacharjee SK. *Handbook of medicinal plant*, Pointer Publication, Jaipur, 2004.p.119-25.
24. Bolla, J.M., Alibert-Franco, S, Handzlik, J., Chevalier, J., Mahamoud, A., Boyer, G., Strategies for bypassing the membrane barrier in multidrug resistant Gram-negative bacteria, *FEBS. Lett.*, 2011; 585(11): 1682-90.
25. Cherry, M. 1985. The needs of the people. In: G.E. Wickens, J.R. Goodin and D.V. Field (ed.). *Plants for arid lands*. Unwin Hyman, London, UK.
26. Cisse, A. A. Ndiaye, P. Lopez-Sall, F. Seck and B. Faye. 2000. Antidiabetic activity of *Ziziphus mauritiana* Lam (Rhamnaceae). *Dakar Med.*, 45(2): 105-107.
27. Cosgrove, S.E., Carmeli, Y., The impact of antimicrobial resistance on health and economic outcomes, *Clin. Infect. Dis.*, 2003; 36(11): 1433-7.
28. D. Senthil Rajan, M. R. (2013). Wound healing activity of an herbal ointment containing the leaf extract of *Ziziphus Mauritiana* Lam. *African Journal of Pharmacy and Pharmacology*, Vol. 7(4), pp. 98-103.
29. Dahiru D, Obidoa O (2008). Evaluation of the antioxidant effects of *Ziziphus mauritiana* lam. Leaf extracts against chronic ethanol-induced hepatotoxicity in rat liver. *Afr. J. Trad. CAM*. 5(1): 3945
30. Dahiru, D., Sini, J.M., John-Africa, L., Antidiarrhoeal activity of *Ziziphus mauritiana* root extract in rodent, *Afr. J. Biotech.*, 2006; 5(10): 941-945. Peto Azam-Ali S, Bonkougou E, Bowe C, Kock D, Godara A, Williams JT. *Ber. International Centre for Underutilised Crops*, Southampton, UK, 2006. p.1-20
31. Dalleau, S., M. Baradat, F. Gueraud and L. Huc. 2013. Cell death and diseases related to oxidative stress: 4-hydroxynonenal (HNE) in the balance. *Cell Death and Differentiation*, 20:1615-1630.
32. Elaloui M, Laamouri A, Fabre J, Mathieu C, Vilarem G, Hasnaoui B. Distribution of free amino acids, polyphenols and sugars of *Ziziphus jujuba* pulps harvested from plants grown in Tunisia. *Natural product research*. 2015;29(1):94-7.
33. Elless, M.P., M.J. Blaylock, J.W. Huang and C.D. Gussman. 2000. Plants as a natural source of concentrated mineral/nutritional supplements. *Food Chemistry*, 71(2): 181-188.
34. Elless, M.P., M.J. Blaylock, J.W. Huang and C.D. Gussman. 2000. Plants as a natural source of concentrated mineral/nutritional supplements. *Food Chemistry*, 71(2): 181-188
35. Gao QH, Wu CS, Wang M. The jujube (*Ziziphus jujuba* Mill.) fruit: a review of current knowledge of fruit composition and health benefits. *Journal of agricultural and food chemistry*. 2013;61(14):3351-63.
36. Gao QH, Wu CS, Wang M. The jujube (*Ziziphus jujuba* Mill.) fruit: a review of current knowledge of fruit composition and health benefits. *Journal of agricultural and food chemistry*. 2013;61(14):3351-63.
37. Grice A.C. (2002) *The Biology of Australian Weeds*. 39. *Ziziphus mauritiana* Lam. *Plant Protection Quarterly*, 17, 2–11.
38. Grosvenor, P.W., Supriono, A., Gray, D.O., *Medicinal plants from Riau Province, Sumatra, Indonesia*. Part 2: antibacterial and antifungal activity, *J. Ethnopharmacol.*, 1995; 45(2): 97-111.
39. Gupta, M.K., A.K. Bhandari and R.K. Singh. 2012. Pharmacognostical evaluation of the leaves of *Ziziphus mauritiana*. *International Journal of Pharmaceutical Sciences and Research*, 3(3): 818-821.
40. Hernandez F, Noguera-Artiaga L, Burlo F, Wojdylo A, Carbonell-Barrachina AA, Legua P. Physico-chemical, nutritional, and volatile composition and sensory profile of Spanish jujube (*Ziziphus jujuba* Mill.) fruits. *Journal of the science of food and agriculture*. 2016;96(8):2682-91.
41. Hocking D. (1993) *Trees for Drylands*. New Delhi, India: Oxford & IBH Publishing. 376pp.
42. Hussain A, Khan MN, Iqbal Z, Sajid MS (2008). An account of the botanical anthelmintics used in traditional veterinary practices in Sahiwal district of Punjab, Pakistan. *J. Ethnopharmacol*. 119: 185– 190.
43. Ivanova, A.S. 1982. Comparative characteristics of the quality composition of some stone fruits and subtropical fruit crops [In Russian] *Bulleten Gosudarstvenno Nikitskogo Botanicheskogo Sada* 47:93-98.
44. Ji X, Peng Q, Yuan Y, Shen J, Xie X, Wang M. Isolation, structures and bioactivities of the polysaccharides from jujube fruit (*Ziziphus jujuba* Mill.): A review. *Food chemistry*. 2017;227: 349-57.
45. Jimoh FO, Adebayo AA, Aliero AA, Afolayan AJ (2008). Polyphenol contents and biological activities of *Rumex ecklonianus*. *Pharm. Biol.*, 45(5): 333-340.

46. Kaaria I (1998). Seed production, dispersal and germination in *Cryptostegia grandifolia* and *Ziziphus mauritiana*, two invasive shrubs in tropical woodlands of Northern Australia. *Australia J. Ecol.*, 21: 324-331
47. Karthikeyan, A., Shanthi, V., Nagasathaya, A., Preliminary phytochemical and antibacterial screening of crude extract of the leaf of *Adhatoda vasica* L., *Int. J. Green Pharm.*, 2009; 3(1): 78-80.
48. Khadri, A.M., S. Neffati, P. Smiti, R.L. Fale, M.L.M. Lino, Serralherio and M.E.M. Araujo. 2010. Antioxidant, antiacetyl cholinesterase and antimicrobial activities of *Cymbopogonschoenanthus* L. Spreng (lemon grass) from Tunisia. *LWT-Food Science and Technology*, 43(2): 331-336.
49. KOLEY, T.K., KAUR, C., NAGAL, S., WALLA, S., JAGGI, S. and SARIKA, J. 2011. Antioxidant activity and phenolic content in genotypes of Indian jujube (*Ziziphus mauritiana* Lamk.). *Arabian J. Chem.* 4, 1–9.
50. Lee Y, Cesario T, Wang Y, Shanbrom E, Thrupp L (2003). Antibacterial activity of vegetables and juices. *Nutr.*, 19: 994-996.
51. Lee, C.K., Kim, H., Moon, K.H., Shin, K.H., Screening and isolation of antibiotic resistance inhibitors from herb materials-resistance inhibition of volatile components of Korean aromatic herbs, *Arch. Pharm. Res.*, 1998; 21(1): 62-6.
52. LI, J., DING, S. and DING, X. 2005. Comparison of antioxidant capacities of extracts from five cultivars of Chinese jujube. *Process Biochem.* 48, 3607–3613.
53. Liu M.J., Zhao Z.H. (2009) Germplasm resources and production of *Ziziphus* in China. *ISHS Acta Horticulturariae*, 840, 25–32.
54. Liu, H., N. Qiu, H. Ding and R. Yao. 2008. Polyphenols contents and antioxidant capacity of 68 Chinese herbals suitable for medicinal or food uses. *Food Research International*, 41(4): 363-370.
55. Lobo, V., A. Patil, A. Phatak and N. Chandra. 2010. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy Review*, 4(8):118-126.
56. Lozoya, M., Lozoya, X., Pharmacological properties in vitro of various extracts of *Mimosa pudica* Linn., *Tepezcohuite Arch. Invest. Mex.*, 1989; 12; 87-93.
57. Lucock, M. 2004. Is folic acid the ultimate functional food component for disease prevention? *BMJ*, 328: 211-214.
58. Marwat, S.K., M.A. Khan, M.A. Khan, F.U. Rehman, M. Ahmad, M. Zafar and S. Sultana. 2009. *Salvadorapersica*, *Tamarixaphylla* and *Ziziphus mauritiana*-three woody plant species mentioned in Holy Quran and Ahadith and their ethnobotanical uses in north western part (D.I. Khan) of Pakistan. *Pakistan Journal of Nutrition*, 8: 542-547.
59. Memon, A.A., N. Memon, D.L. Luthria, A.A. Pitafi and M.I. Bhangar. 2012a. Phenolic compounds and seed oil composition of *Ziziphus mauritiana* L. *Fruit. Polish Journal of Food and Nutrition Sciences*, 62(1): 15-21
60. Mesaik AM, Poh HW, Bin OY, Elawad I, Alsayed B. In Vivo Anti-Inflammatory, AntiBacterial and Anti-Diarrhoeal Activity of *Ziziphus Jujuba* Fruit Extract. Open access Macedonian journal of medical sciences. 2018;6(5):757-66.
61. Michel A (2002). Tree, Shrub and Liana of West African Zone. Margraf Publishers GMBH, Paris, pp. 440.
62. Mishra, T., M. Khullar and A. Bhati. 2011. Anticancer potential of aqueous ethanol seed extract of *Ziziphus mauritiana* against cancer cell lines and ehrlich ascites carcinoma. *Evidence Based Complementary and Alternative Medicine*, 2011: 1-11.
63. miyasaki, Y., J.D. Rabenstein, J. Rhea, M.L. Crouch, U.M. Mocek, P.E. Kittell, M.A. Morgan, W.S. Nichols, M.M.V. Benschoten, W. D. Hardy and G.Y. Liu. 2013. Isolation and characterization of antimicrobial compounds in plant extracts against multidrug-resistant *Acinetobacter baumannii*. *PLoS One*, 22: 1-8.
64. Morton J.F. (1987) Indian jujube. In *Fruits of Warm Climates*, pp. 272–275. Miami, FL, USA: Florida Flair Books.
65. Morton, J., Indian Jujube. In: *Fruits of warm climates*. Julia, F., Morton, J. (eds) Miami, FL., 1987; 272275.
66. Morton, J., Indian Jujube. In: *Fruits of warm climates*. Julia, F., Morton, J. (eds) Miami, FL., 1987; 272275.
67. N Sunitha*, K Swathi, N Rama Rao, Method development and validation of quercetin obtained from the extraction of almond leaves by UV spectrophotometry and FTIR, *international Journal of botany study*, 2021.
68. Najafi, S., Sanadgol, N., Sadeghinejad, B., Ashofteh Beira, S., Phytochemical screening and antibacterial activity of *Citrullus Colocynthis* (L.) Schrad. against *Staphylococcus aureus*, *Journ. Oral Med. Path. Radio.*, 2010; 4(9): 2321-2325.
69. Nascimento, G.G.F., Locatelli, J., Freitas, P.C., Silva, GL. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria, *Braz. J. Microbiol.*, 2000; 31(4): 247-56.
70. Okoko T, Oruambo IF (2008). The effects of *Hibiscus sabdariffa* calyx on cisplatin-induced tissues damaged in rats. *Biochem. Streak*, 20(2): 47-52.
71. Pareek OP. *Fruits for the Future 2: Ber*, International Centre for Underutilised Crops, University of Southampton, Southampton, UK, 2001.p.1-32.
72. Pareek S., Yahia E.M. (2013) Postharvest biology and technology of Ber fruit. In *Horticultural Reviews*. Volume 41, pp. 201–240. Ed. J. Janick. Hoboken, NJ, USA: John Wiley & Sons.
73. Parmar, P., Bhatt, S., Dhyani, S., Jain, A., Phytochemical studies of the secondary metabolites of *Ziziphus mauritiana* Lam. *Leaves, Int. J. Pharma. R.*, 2012; 4(3): 153-155.
74. Peto Azam-Ali S, Bonkoungou E, Bowe C, Kock D, Godara A, Williams JT. *Ber*, International Centre for Underutilised Crops, Southampton, UK, 2006. p.1-20

75. Qureshi, K.A., Bholay, A.D., Elhassan, G.O, Khan, R.A., El-Agamy, E.I., Isolation, purification and characterization of bacteriocin produced by *Bacillus pumilus* NJ-M2; A Future Biopreservative, Intern. Journ. Biol. Pharm. Allied Scien., 2016; 5(11): 2840-2862.
76. Senthil Kumar M, Sripriya R, Vijaya Raghavan H, Sehgal P (2006). Wound Healing Potential of *Cassia fistula* on Infected Albino Rat Model. *J. Surg. Res.* 131: 283-289.
77. Shahidi, F. 2009. Nutraceuticals and functional foods: Whole versus processed foods. *Trends in Food Science and Technology*, 20(9):376-387.
78. Siddharth, P., P. Kailash, V. Niraj, M. Karuna, P. Vimal, P. Bharadia and K. Pundarikakshudu. 2010. Antiulcer activity of methanolic extract of *Ziziphus mauritiana* stem bark. *International Journal of Pharmacognosy and Phytochemical Research*, 2(3): 6-11.
79. Sobhani Z, Nikoofal-Sahlabadi S, Amiri MS, Ramezani M, Emami SA, Sahebkar A. Therapeutic Effects of *Ziziphus jujuba* Mill. Fruit in Traditional and Modern Medicine: A Review. *Medicinal chemistry*. 2020;16(8): 1069-88.
80. Sofowora, A., *Medicinal Plants and Traditional medicine in Africa*. Published by John Wiley and Sons Ltd. 1st edition., 1982; 131: 168-171.
81. Stadlmayr B, Charrondiere UR, Eisenwagen S, Jamnadass R, Kehlenbeck K. Nutrient composition of selected indigenous fruits from sub-Saharan Africa. *Journal of the science of food and agriculture*. 2013;93(11):2627-36.
82. Szeto YP, Tomlinson B, Benzie IFF (2002). Total antioxidant and ascorbic acid content of fresh fruits and vegetables: Implications for dietary planning and food preservation. *Brit. J. Nutr.*, 87: 55-59.
83. Tahergorabi Z, Abedini MR, Mitra M, Fard MH, Beydokhti H. "Ziziphus jujuba": A red fruit with promising anticancer activities. *Pharmacognosy reviews*. 2015;9(18): 99-106.
84. Wang, S.Y., C.Y. Tian, Y.Z. Ye, Q.S. Yeng, Y.Z. Wang, Q.L. Zhang and L.M. Fan. 1992. The wild fruit germplasm resources of Hanan Province. *J. Fruit Sci.* 19:239-242.
85. Zeyadi M, Almulaiky YQ. A novel peroxidase from *Ziziphus jujuba* fruit: purification, thermodynamics and biochemical characterization properties. *Scientific reports*. 2020;10(1):8007.